WORLD INTELLECTUAL PROPERTY ORGANIZATION



ATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

92166853

(11) International Publication Number:

WO 92/06675

ΑI

(43) International Publication Date:

30 April 1992 (30.04.92)

(21) International Application Number:

PCT/US91/07574

(22) International Filing Date:

9 October 1991 (09.10.91)

(30) Priority data:

599,694

18 October 1990 (18,10,90)

pean patent), CA, CH (European Fatent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European pa-

(81) Designated States: AT (European patent), AU, BE (Euro-

tent).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of

amendments.

(71) Applicant: MINNESOTA MINING AND MANUFAC-1 TURING COMPANY JUS USJ: 3M Center, Post Office Box 33427, Saint Paul, MN 55133-3427 (US).

(72) Inventors: SCHULTZ, Robert, K.; SCHULTZ, David, W. : Post Office Box 33427, Saint Paul, SSS 53133-3427 (US).

(74) Agents: REEDICH, Douglas, E. et al.; Intellectual Property Counsel, Minnesota Mining and Manufacturing Company, Post Office Box 33427, Saint Paul, MN 55133-3427 (US)

(54) Title: AEROSOL FORMULATION COMPRISING BECLOMETHASONE 17,21 DIPROPIONATE

(57) Abstract

THE PARTY OF THE PROPERTY OF THE PARTY OF TH

Pharmaceutical solution acrosol formulations comprising beclomethasone 17,21 dipropionate, ethanol, and a propellant selected from the group consisting of 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, and a mixtuse thereof-

AGRESOL FORMULATION COMPRISING BECLOMETHASONE 17,21-DIPROPIONATE T

TREATING BRONCHAL ASTAMA

BUCCAL OF PULMONARY,

Phanol as cosolvent => Solution system.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	E.S	Spain	MC	Madagascar
AU	Australia	FI	Finland	ML	Mali
88	Bartonios	FR.	France	MN	Monzolu
33	Belgium	CA	Guñon	MR	Mauritania
6.F	Burking Favo	CB	United Kingdom	MW	Malawi
DC.	Bulgaria	CH	Guinea	NL.	Netherlands
£ 7	Benin	CR	Grace	NO	Norman
BR	Brazil	HU	Hungary	FL	Poland
C۸	Canada	17	lul,	80	Romania
CF	Central African Republic	٦٢.	Lipin	a2	Sutin
CC	Congo	KP	Democratic People's Republic	32	Sweden
CS	Switzerland		of Koria	SN	Semegat
CI	Côte d'Ivoire	KX	Republic of Kores	SU*	Server Union
CH	Camarain	LI	Lechtenstein	το	Chad
C.	Czechoslovakiu	LA.	Sri Lanka	TC	Toes
DE.	Germany	LU	Lusembourg	US	United States of America
.			· · · · · · · · · · · · · · · · · ·	-	

THE PROPERTY OF THE PROPERTY O

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

Aerosol formulation comprising beclomethasone 17,21 dipropionate.

TECHNICAL FIELD OF THE INVENTION

This invention pertains to solution aerosol formulations suitable for use in administering drugs. In another aspect this invention pertains to formulations comprising beclomethasone 17,21 dipropionate.

10

20

25

5

BACKGROUND OF THE INVENTION

Pharmaceutical suspension aerosol formulations currently use a mixture of liquid chlorofluorocarbons as the propellant. Fluorotrichloromethane,

dichlorodifluoromethane and dichlorotetrafluoroethane are the most commonly used propellants in aerosol formulations for administration by inhalation.

Chlorofluorocarbons have been implicated in the destruction of the ozone layer and their production is being phased out. Hydrofluorocarbon 134a (EFC-134a, 1,1,1,2-tetrafluoroethane) and hydrofluorocarbon 227 (EFC-227, 1,1,1,2,3,3,3-heptafluoropropane) are viewed as being less destructive to ozone than many chlorofluorocarbon propellants; furthermore, they have low toxicity and vapor pressure suitable for use in aerosols.

Beclomethasone 17,21 dipropionate is commercially available as an aerosol product comprising a suspension of a chlorofluorohydrocarbon solvate of beclomethasone 17,21 dipropionate in

- chlorofluorohydrocarbon propellants. Preparation of the solvate requires several processing steps and is required in order to obtain a stable aerosol formulation, i.e., one in which the micronized particles of active ingredient remain in the desired respirable particle size range. A
- solution formulation of beclomethasone 17,21 dipropionate could simplify formulation manufacture and increase the respirable fraction (i.e., the percentage of active ingredient able to reach the airways of the lung where the pharmaceutical effect is exerted).

10

15

20

25

30

35

40

U.S. Pat. No. 2,868,691 discloses a self-propelling pharmaceutical aerosol formulation comprising i) a medicament; ii) a propellant represented generally by the formula $C_mH_mCl_\gamma F_z$, wherein m is an integer less than 3, n is an integer or zero, y is an integer or zero, and z is an integer, such that n + y + z = 2m + 2; and iii) a cosolvent which assists in the dissolution of the medicament in the propellant. Ethanol is an example of a cosolvent disclosed in this patent. The above formula representing the propellant component generically embraces HFC-134a. This patent does not, however, disclose beclomethasone 17,21 dipropionate or suggest how stable solution aerosol formulations (i.e., formulations that are chemically stable and exhibit desirable respirable fraction) containing any propellant and beclomethasone 17,21 dipropionate can be prepared.

SUMMARY OF THE INVENTION

The present invention provides an aerosol formulation comprising a therapeutically effective amount of beclomethasone 17,21 dipropionate, a propellant comprising a hydrofluorocarbon selected from the group consisting of 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, and a mixture thereof, and ethanol in an amount effective to solubilize the beclomethasone 17,21 dipropi ate in the propellant, the formulation being further characterized in that substantially all of the beclomethasone 17,21 dipropionate is dissolved in the formulation, and the formulation is substantially free of any surfactant.

Certain of the preferred formulations of the invention exhibit very desirable chemical stability and provide respirable fractions significantly greater than commercially available beclomethasone 17,21 dipropionate products. Moreover, the formulations of the invention are convenient to manufacture since no solvate of the active ingredient need be prepared.

The pharmaceutical solution aerosol formulations of the invention are suitable for pulmonary, buccal, or nasal administration.

10

20

25

30

DETAILED DESCRIPTION OF THE INVENTION

All weight percentages recited herein are based on the total weight of the formulation unless otherwise indicated.

The medicament beclomethasone 17,21 dipropionate is generally present in a formulation of the invention in a therapeutically effective amount, i.e., an amount such that one or more metered volumes of the formulation contains an amount of drug effective to exert the intended therapeutic action. Preferably the medicament will constitute about 0.02 to about 0.6 percent by weight, more preferably about 0.05 to about 0.5 percent by weight of the total weight of the formulation.

Ethanol is generally present in an amount effective to solubilize the beclomethasone 17,21 dipropionate in the propellant. Preferably, ethanol constitutes about 1 to about 20 percent by weight of the total weight of the aerosol formulation. More preferably, ethanol constitutes about 2 to about 12 percent by weight and even more preferably about 2 to about 10 percent by weight of the aerosol formulation. Most preferably, ethanol will be present in an amount sufficient to dissolve substantially all of the medicament present in the formulation and to maintain the medicament dissolved over the time period and conditions experienced by commercial serosol products, but not substantially in excess of said amount. Particularly desirable formulations of the invention, while not containing amounts of ethanol substantially in excess of that required (during manufacture of the formulation) to dissolve the amount of active ingredient employed, may be subjected to a temperature of -20°C. without precipitation of the active ingredient.

The hydrofluorocarbon propellant can be HFC134a, HFC-227, or a mixture thereof. The propellant
preferably constitutes from about 50 to about 99 percent by
weight, preferably from about 58 to about 98 percent by
weight, and more preferably about 90 to about 98 percent by
weight of the total weight of the aerosol formulation. The
hydrofluorocarbon propellant is preferably the only

15

20

30

propellant present in the formulations of the invention. However, one or more other propellants (e.g., 1-chloro-1,1-difluoroethane) can also be present.

The formulations of the invention are substantially free of any surfactant. By "substantially free" as used in the instant specification and claims is meant that the formulations contain no more than 0.0005 percent by weight of a surfactant based on the total weight of the formulation. Preferred formulations contain no surfactant. Presence of a significant amount of a surfactant is believed to be undesirable in the case of solution formulations of beclomethasone 17,21 dipropionate because surfactants such as oleic acid and lecithin seem to promote chemical degradation of the active ingredient when the latter is dissolved in the mixture of HFC-134a and ethanol.


preferred formulations according to the invention consist essentially of beclomethasone 17,21 dipropionate in an amount of about 0.05 to about 0.35 percent by weight based on the weight of the total formulation, ethanol in an amount of about 2 to about 8 percent by weight based on the total weight of the formulation, and 1,1,1,2-tetrafluoroethane.

The solution formulations of the invention can be prepared by dissolving the desired amount of beclomethasone 17,21 dipropionate in the desired amount of anhydrous ethanol accompanied by stirring or sonication. The aerosol vial may then be filled using conventional cold-fill or pressure-fill methods.

The following examples are provided to illustrate the invention but should not be construed as limiting the invention.

Examples 1 - 7

ingredients (TABLE I) in the indicated amounts were prepared with the percentages being expressed in parts by weight based upon the total weight of the particular formulation. The active ingredient employed in preparing the formulations of Examples 2, 3, and 5 - 7 was



15

20

beclomethasone dipropionate, USP while that employed in preparing the formulations of Examples 1 and 4 was a conventional trichloromonofluoromethane solvate of beclomethasone dipropionate. The formulations of Examples 1, 4, 5 and 6 were prepared by i) dissolving the active ingredient in the ethanol; ii) metering the solution obtained above into an aluminum vial and crimping a continuous valve onto the vial; iii) pressure-filling the vial with 1,1,1,2-tetrafluoroethane; iv) chilling the vial to -60°C.; and v) replacing the continuous valve with a 50 microliter valve which is available under the trade designation "W303-98" from 3M. The formulations of Examples 2, 3 and 7 were prepared by i) dissolving the active ingredient in the ethanol; ii) metering the solution obtained above into an aluminum vial and crimping a 50 microliter pressure-fill valve which is available under the trade designation Spraymiser M3652 from 3M onto the vial; and iii) pressure-filling the vial with 1,1,1,2tetrafluoroethane.

The actuator employed in the case of all the formulations was a solution actuator available under the trade designation "M3756" from 3M. The elastomer employed in the valves in the case of all formulations was that available under the trade designation "DB-218" from American Gasket and Rubber Co. (Chicago, IL.) 25

TABLE I

Ingredient		Example					
	1	2	3	4	5	6	7
Beclomethasone 17,21 Dipropionate	0.1%	0.1%	0.25%	0.3%	0.4%	0.44%	0.5%
Ethanol (anhydrous)	3%	5%	10%	5%	10%	10%	15%
1,1,1,2- Tetra- fluoroethane	96.9%	94.9%	89.75%	94.7%	89.6%	89.56%	84.5%

The chemical stability of the formulation of Example 4 was determined in respect to recovery of the active ingredient over time when the formulation was stored at 40°C. TABLE II contains the data.

TABLE II

					····	
10	Storage Time (Weeks)	0	2	4	7	12 ,
	<pre>% Recovery</pre>	101.4, 98.7	101.9, 101.6	100.8, 99.6	99.3, 95.5	100.6

15

5

The formulation of Example 1 did not exhibit precipitation of the active ingredient on freezing to -60°C.

The respirable fraction provided by the formulations of Examples 1 - 7 was determined using an Anderson MK II Cascade Impactor with the average respirable fraction obtained from each being in excess of 40%. In the case of the formulations of Examples 1 and 4, the respirable fraction was about 76% and about 70%, respectively.

25

20

From the above data, it is believed that the optimum amount of active ingredient for low and high strength products would be about 0.08 and 0.34 percent by

10.

20

25

3.0

35

*weight, respectively, based on the total weight of the formulations.

Example 8

A mixture containing 1.67 g of beclomethasone 17,21 dipropionate and 160g of cold (-65°C) ethanol was homogenized using a Virtis 45 homogenizer. The resulting suspension was placed in a one gallon stainless steel filling vessel equipped with a stir bar. A 1839 g portion of cold (-65°C) 1,1,1,2-tetrafluoroethane was added to the filling vessel. After about 5 minutes of stirring, a solution was obtained. The resulting formulation contained 0.08 percent by weight of beclomethasone 17,21 dipropionate, 8.0 percent by weight of ethanol and 91.92 percent by weight of 1,1,1,2-tetrafluoroethane. 15 formulation was cold filled into aerosol vials and then 50 μL cold fill valves were crimped onto the vials.

Example 9

Using the general method of Example 8, a formulation containing 0.34 percent by weight of beclomethasone 17,21 dipropionate, 8.0 percent by weight of ethanol and 91.66 percent by weight of 1,1,1,2tetrafluoroethane was prepared. The formulation was cold filled as a suspension into aerosol vials which were then equipped with 50 μ L cold fill valves. The formulation changed from a suspension to a solution as the vials warmed to room temperature.

Example 10

A formulation containing 0.3 percent by weight of beclomethasone 17,21 dipropionate, 10 percent by weight of ethanol and 89.7 percent by weight of 1,1,1,2,3,3,3heptafluoropropane was prepared by i) weighing a 30 mg portion of beclomethasone 17,21 dipropionate into an aerosol vial ii) crimping a continuous valve onto the vial and iii) pressure filling with a solution containing 10 percent ethanol in 1,1,1,2,3,3,3-heptafluoropropane.

25

WHAT IS CLAIMED IS:

- therapeutically effective amount of beclomethasone 17.21
 dipropionate, a propellant comprising a hydrofluorocarbon selected from the group consisting of 1.1.1.2tetrafluoroethane, 1.1.1.2.3.3.3-heptafluoropropane, and a mixture thereof, and ethanol in an amount effective to solubilize the beclomethasone 17.21 dipropionate in the propellant, the formulation being further characterized in that substantially all of the beclomethasone 17.21 dipropionate is dissolved in the formulation, and that the formulation is substantially free of any surfactant.
- 2. A solution aerosol formulation according to Claim 1, comprising between 0.02 and about 0.6 percent by weight beclomethasone 17,21 dipropionate, between about 1 and about 20 percent by weight ethanol, and between about 80 and about 99 percent by weight of said propellant.
 - 3. A solution aerosol formulation according to Claim 1 wherein said beclomethasone 17,21 dipropionate is present in an amount of about 0.05 to about 0.5 percent by weight.
 - 4. A solution aerosol formulation according to Claim 1 wherein said ethanol is present in an amount of about 2 to about 12 percent by weight.
- 5. A solution aerosol formulation according to Claim 1 wherein said ethanol is present in an amount of about 2 to 10 percent by weight.
- 6. A solution aerosol formulation according to Claim 1 wherein said propellant is present in an amount of about 88 to about 98 percent by weight.

15

20

25

30

35

- 7. A solution aerosol formulation according to Claim 1 comprising 1,1,1,2-tetrafluoroethane as substantially the only propellant.
- 8. A solution aerosol formulation according to Claim 1 comprising 1,1,1,2,3,3,3-heptafluoropropane as substantially the only propellant.
- 9. A solution aerosol formulation according to Claim 1 comprising beclomethasone 17,21 dipropionate in an amount of about 0.05 to about 0.5 percent by weight, ethanol in an amount of about 2 to about 12 percent by weight and said propellant in an amount of about 88 to about 98 percent by weight.
 - 10. A solution aerosol formulation according to Claim 1 comprising beclomethasone 17, 21 dipropionate in an amount of about 0.05 to about 0.45 percent by weight, ethanol in an amount of about 2 to about 10 percent by weight and said propellant in an amount of about 90 to about 98 percent by weight.
 - 11. A solution aerosol formulation according to Claim 1, consisting essentially of beclomethasone 17,21 dipropionate in an amount of about 0.05 to about 0.35 percent by weight, ethanol in an amount of about 2 to about 8 percent by weight, and 1,1,1,2-tetrafluoroethane.
 - 12. A solution aerosol formulation according to Claim 1, wherein the amount of ethanol present is not substantially in excess of the amount required to dissolve substantially all of the beclomethasone 17,21 dipropionate, but is sufficient to permit said formulation to be subjected to a temperature of -20°C. Without significant precipitation of said beclomethasone 17, 21 dipropionate.
 - 13. A method of treating bronchial asthma in a mammal comprising administering to said mammal an amount of

a formulation according to Claim 1 sufficient to treat the asthmatic condition.

14. A method of preparing a solution aerosol formulation comprising the step of combining a therapeutically effective amount of beclomethasone 17,21 dipropionate, a propellant selected from the group consisting of 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, and a mixture thereof, and an amount of ethanol effective to solubilize the beclomethasone 17,21 dipropionate in the propellant.

INTERNATIONAL SEARCH REPORT

• .	int	ernational Application No PCT/U	S 91/07574
SSIFICATION		a symbols apply, indicate all)	
raing to Internat	tional Patent Classification (IPC) or to cour restor	at Classification and IPC	
έ: Λ 61 K	9/12		
FIELDS SEARCH	ED	- Seambed	
	Minimum Documentation	ification Symbols	
eilication System	<u> </u>	The 2110th Dynamic Control of the 1	
:5	A 61 K		
.5		n Minimum Documentation	į
	Documentation Searched binds the to the Extent that such Documents ar	e Included in Fields Searched	
	CONSIDERED TO BE RELEVANT		
	ition of Document, 11 with Indication, where approp	priate, of the relevant passages 12	Relevant to Claim Kc.13
cita	1, 9111173 (FISONS PLC) 8 Au	gust 1991,	1-14
(WO, A	see the whole document	•	
	and the second of the second o		
	A1, 9111495 (BOEHRINGER INGEL	HEIM	1-14
X 140, 4	INTERNATIONAL) 8 August 1991	•	
	see the whole document		
	see one more as		
			
	OUTACOUTA LABORATION	TES THE	1-14
WO,	A1, 8603750 (RIKER LABORATOR	11.5, 11.01,	
	3 July 1986, see the whole document		
	See the whole Toland		
		•	
. 1			•
		•	1.7
	·		
			'
		bliched al	ner the international filing da
* Special cale	egories of cited documents: 18	or priority date and not in c grided to understand the prin	ciple or theory underlying the
"A" document	defining the central state of the	Invention	as a section of Imposition
ezdier de	cument but published on or area. The	CENTINE BU INVENTIVE STOS	
ביי בטכעות אול	which may throw doubts on priority claim(s) or	ove document of particular rele	vance, the claimed invention plye an inventive step when to come or more other such some
ritation 0	r other special reason (as specifical	secument is combined with	end opyions to a betach accurate to a betach accura
"C" document	t reforming to an oral disclosure, use, exhibition o	in me ar-	
	ans t published prior to the international filing date b n the priority date claimed	ut "8" cocument member of the s	
ISTER MAI	TION	Date of Mailing of this Internation	nat Swarch Report
Date of the Actu	nt Completion of the International Saurch	2 5. 63. 92	•
9th March	•		
		Signature of Authorized Officer	D. (M
International Se	UROPEAN PATENT OFFICE	Mme. M. van dar	Drift
F	OKOREVU AVIEMI OLITOR	Willer Mr. 4411. 00.	

The Artist (second speed (January 1985)

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/US 91/07574

SA

54160

annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report members are as contained in the European Patent Office EDP file on 01/02/92 European Patent office is in no way fiable for these particulars which are merely given for the purpose of information.

Psient document cited in search report	Publication date	Patent mem	Publication date	
)-Al- 9111173	08/08/91	NONE		
)-A1- 9111495	08/08/91	AU-D- DE-A-	7211391 4003272	21/08/91 08/08/91
)-A1- 8603750	03/07/86	AU-B- AU-D- CA-A- EP-A-B- JP-T- US-A-	587010 5308786 1253806 0205530 62501706 4810488	03/08/89 22/07/86 09/05/89 30/12/86 09/07/87 07/03/89

For more details about this annex : see Official Journal of the European patent Office, No. 12/92

FPO FORM FOATE